#### IMPORTANT INFORMATION

#### FOR USE ONLY AS A PROFESSIONAL HERBICIDE

Crops / Situations:
Maximum Individual Dose:
Maximum number of applications
Maximum Total Dose:
Latest time of application

Amenity grassland, non-crop areas 200 mL per 10 litres water or 4l/ha 1 per vear 4.0 litres product per hectare per vear 31<sup>st</sup> August

DO NOT USE between 31st August and 1st March.

For applications made between 1<sup>st</sup> March and 31<sup>st</sup> May, 1 application may be made per calendar

For applications made between 1<sup>st</sup> June and 31<sup>st</sup> July, only 1 application may be made in a 2 year period

For applications made between 1st August and 31st August, only 1 application may be made in a 3 year period.


DO NOT USE on land that will be grazed by livestock.

DO NOT USE on land where vegetation will be cut for animal feed, fodder or bedding nor for composting or mulching within one calendar year of treatment.

DO NOT USE on airfields.

DO NOT apply spot applications to more than 20% of the area. Users must have received adequate instruction, training and guidance in the safe use of the product and must take all reasonable precautions to protect the health of human beings, creatures and plants and safeguard the environment.

Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Follow the Code of Practice for Using Plant Protection Products.





# Garlon<sup>®</sup> Ultra

#### HERBICIDE

<sup>®</sup> Trademarks of Corteva Agriscience and its affiliated companies.



#### Product Registration Number: PCS No. 05396

A soluble concentrate containing 12 g ae/litre aminopyralid and 120 g ae/litre triclopyr.

For the control of certain PERENNIAL, HERBACEOUS and WOODY WEEDS in amenity grassland.

#### Authorisation Holder:

#### Corteva Agriscience UK Limited

CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE, United Kingdom Telephone: +44 (0) 1462 457272 24 Hour Emergency Telephone Number: +353(0) 76 680 5288 Technical Hotline: +44 (0) 800 689 8899, e-mail: ukhotline@corteva.com National Poisons Centre: 01 809 2166

Marketing Company Whelehan Crop Protection Suite 12 Bunkilla Plaza, Bracetown Business Park Clonee, Ireland, D15 WR13 Tel: + 353 (1) 574 2410

(H)

## SAFETY PRECAUTIONS

#### Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION

(FACESHIELD) when handling the concentrate. WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when applying by hand-held

equipment.

WHEN USING DO NOT EAT, DRINK OR SMOKE. WASH CONCENTRATE from skin or eves immediately.

#### Environmental protection:

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area. DO NOT CONTAMINATE WATER with the product or its container. (Do not clean application equipment near surface water / Avoid contamination via drains from farmvards and roads.)

#### Storage and disposal:

KEEP OUT OF REACH OF CHILDREN. KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS. KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place, **RINSE CONTAINER THOROUGHLY** by using an integrated pressure rinsing device or manually rinsing three times. Add washings to 2310 spraver at time of filling and dispose of safely.

#### READ DIRECTIONS FOR USE ON ATTACHED LEAFLET PROTECT FROM FROST

FOR PROFESSIONAL USE ONLY

Triple Rinse Containers, Puncture And Invert To Dry At Time Of 🔊 🔊 Use

> Artwork Free area for in-house print of 2D data matrix code



NET CONTENTS : 1 LITRE 🕀

## **DIRECTIONS FOR USE**

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

#### WARNINGS

Take extreme care to avoid drift onto susceptible crops, non-target plants or waterways. Do not apply directly to, or allow spray drift to come into contact with agricultural or horticultural crops, amenity plantings, gardens, ponds, lakes or watercourses.

**SENSITIVE PLANTS** may be harmed by residues of GARLON ULTRA in soil and treated vegetation. Do not apply GARLON ULTRA on or adjacent to soil which may be used as garden top-soil, potting soil, etc. or to grass which may be cut and used as mulch or for compost for horticultural or garden crops. Do not use cuttings from treated grass for mulching or composting.

Cut treated grass must not be removed from site, or used for animal feed, animal bedding, composting or mulching

Treated grass must not be used for grazing.

On level ground there is negligible lateral movement but do not apply GARLON ULTRA around desirable trees or shrubs, do not spray under the canopy and 1 metre at least from the trunk of desirable trees or shrubs. Care should be taken on slopes to prevent leaching into areas where desirable shrubs, etc are present.

#### NOTES

Grass and weeds must be actively growing to ensure good weed control and minimal check to the grass. Therefore do not spray in drought, hot or very cold weather conditions.

Do not use on grass less than one year old.

To allow maximum translocation to the roots do not cut grass for 7 days after application.

Clover will be killed by application of GARLON ULTRA.

Wash equipment thoroughly with water and detergent immediately after use.

Non-printed area on page 2 ONLY (on the back of 2D code area printed on the front page)

#### HERBACEOUS WEED CONTROL

#### BROADCAST APPLICATION WEEDS CONTROLLED, RATES OF USE AND TIMING OF APPLICATION

Weeds	Rates of use litres/ha	Optimum timing of application
Bramble <sup>1</sup>	4.0	Treat when the bramble is actively growing but is less than 50 cm high.
Common mugwort	4.0	Treat when the mugwort is actively growing and less than 70 cm high.
Common nettle <sup>1</sup>	2.0	Treat when the nettles are actively growing and less than 70 cm high.
Creeping thistle	4.0	Treat when the thistle is actively growing and less than 70 cm high.
Hogwe <b>ed</b> <sup>2</sup>	4.0	Treat when the hogweed is actively growing and less than 70 cm high.
Rosebay willowherb	4.0	Treat when the willowherb is actively growing and less than 15 cm high.

<sup>1</sup> A second application may be required the following year.

<sup>2</sup> Moderately susceptible

#### **WOODY AND HERBACEOUS WEED CONTROL**

### LOCALISED SPOT APPLICATION

#### WEEDS CONTROLLED, RATES OF USE AND TIMING OF APPLICATION

Weeds	Rate of use mL/10 litres of water	Optimum timing of application
Bramble	150	Up to 1 metre high.
Broom	200	Up to 1 metre high.
Buddleia	200	Up to 1 metre high.
Gorse	150	Up to 1 metre high.
Japanese knotweed	200	1 metre high, with good foliage cover.

#### TIMING OF APPLICATION

GARLON ULTRA should be applied between March and the end of August. The timing of application of GARLON ULTRA is crucial and for good results GARLON ULTRA must be applied to actively growing weeds.

#### APPLICATION

#### Broadcast Treatment

GARLON ULTRA should be applied through a tractor-mounted hydraulic sprayer provided it is in good working order and has been calibrated according to the manufacturers' recommendations.

#### Mixing

Fill the spray tank half full with water and add the required amount of GARLON ULTRA mixing well. Top up with water and continue agitation until the spray tank is full. Maintain agitation while spraying. Use the spray immediately.

#### Spray Volume

For overall application GARLON ULTRA should be used in a spray volume between 300L and 600L per hectare to give good coverage of the weeds. Higher water volumes are recommended where weed densities are high and where the grass is dense at the time of application.

#### Spray Quality

Apply as a MEDIUM quality spray as defined by the BCPC system.

#### Spot Treatment

For localised treatment using a suitable lance from a knapsack or tractor mounted sprayer, use a solution of 150-200 mL of GARLON ULTRA depending on the target weed, per 10 litres of water. Woody weeds should be thoroughly wetted with the spray solution, but spraying until «run-off» will decrease activity. The use of flood jets is recommended to prevent drift. Care should be taken to avoid local overdosing. Woody weeds should not exceed a 1 metre in height.

#### TRADEMARK ACKNOWLEDGEMENTS

GARLON ULTRA is a trademark of Corteva Agriscience and its affiliated companies. All other brand names are trademarks of other manufacturers for which proprietary rights may exist.

#### **Corteva Agriscience Conditions of Supply**

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

#### To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



https://www.corteva.ie/content/dam/dpagco/corteva/eu/ie/en/files/sds/ GARLON-ULTRA-IRL-SDS.pdf

Alternatively contact your supplier

# THIS PAGE HAS BEEN LEFT BLANK INTENTIONALLY

#### IMPORTANT INFORMATION

#### FOR USE ONLY AS A PROFESSIONAL HERBICIDE

Crops / Situations:	Amenity of
Maximum Individual Dose:	200 mL p
Maximum number of applications:	1 per vea
Maximum Total Dose:	4.0 litres
Latest time of application	31 <sup>st</sup> Augu

grassland, non-crop areas per 10 litres water or 4l/ha product per hectare per vear

DO NOT USE between 31st August and 1st March.

For applications made between 1<sup>st</sup> March and 31<sup>st</sup> May, 1 application may be made per calendar

For applications made between 1st June and 31st July, only 1 application may be made in a 2 year period

For applications made between 1st August and 31st August, only 1 application may be made in a 3 year period

DO NOT USE on land that will be grazed by livestock.

DO NOT USE on land where vegetation will be cut for animal feed, fodder or bedding nor for composting or mulching within one calendar year of treatment.

DO NOT USE on airfields.

DO NOT apply spot applications to more than 20% of the area. Users must have received adequate instruction, training and guidance in the safe use of the product and must take all reasonable precautions to protect the health of human beings, creatures and plants and safeguard the environment.

Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Follow the Code of Practice for Using Plant Protection Products.



SAFETY INFORMATION

Garlon Ultra® UFI: AN65-D0D0-M00C-NOHC

#### WARNING

**Causes serious eve irritation** May cause damage to kidneys through prolonged or repeated exposure Very toxic to aquatic life with long lasting effects

Wash hands thoroughly after handling. Wear protective gloves / protective clothing / eye protection / face protection. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do - continue rinsing. If eye irritation persists: get medical advice / attention. Collect spillage. Contains Triclopyr. May produce an allergic reaction. Dispose of contents / container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. 05396



## Garlon<sup>®</sup> Ultra

#### **HERBICIDE**

<sup>®</sup> Trademarks of Corteva Agriscience and its affiliated companies.



HERBICIDES

#### Product Registration Number: PCS No. 05396

A soluble concentrate containing 12 g ae/litre aminopyralid and 120 g ae/litre triclopyr.

For the control of certain PERENNIAL, HERBACEOUS and WOODY WEEDS in amenity grassland.

#### Authorisation Holder:

#### Corteva Agriscience UK Limited

CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE, United Kingdom Telephone: +44 (0) 1462 457272 24 Hour Emergency Telephone Number: +353(0) 76 680 5288 Technical Hotline: +44 (0) 800 689 8899, e-mail: ukhotline@corteva.com National Poisons Centre: 01 809 2166

Marketing Company Whelehan Crop Protection Suite 12 Bunkilla Plaza, Bracetown Business Park Clonee, Ireland, D15 WR13 Tel: + 353 (1) 574 2410

#### NET CONTENTS : 1 LITRE 🕀



#### Operator protection:

 $(\mathbf{H})$ 

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION

(FACESHIELD) when handling the concentrate. WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND

SUITABLE PROTECTIVE GLOVES when applying by hand-held equipment.

WHEN USING DO NOT EAT, DRINK OR SMOKE. WASH CONCENTRATE from skin or eves immediately.

#### Environmental protection:

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

DO NOT CONTAMINATE WATER with the product or its container. (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads.)

#### Storage and disposal:

KEEP OUT OF REACH OF CHILDREN. KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS. KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place. RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to spraver at time of filling and dispose of safely.

#### READ DIRECTIONS FOR USE ON ATTACHED LEAFLET PROTECT FROM FROST

FOR PROFESSIONAL USE ONLY

Triple Rinse Containers, Puncture And Invert To Dry At Time Of Use