

Technical Information Sheet

IRELAND

FUNGICIDE

Aquino™ is a new fungicide for the control of *Septoria tritici*, brown rust and yellow rust in Winter Wheat, Winter Triticale, Winter Spelt and Durum Wheat.

Aquino is broad-spectrum fungicide containing Fenpicoxamid, more commonly known as Inatreq Active. Aquino is part of a new class of chemistry called picolinamides which has a new site of action against fungi in cereals. Aquino will be supplied with Turret® 60 as a co-pack

Key facts

Product Registration Number:	Aquino – PCS 06751 UFI code: 9059-M0XQ-2002-81QX Turret 60 – PCS 06384		
Active Ingredient:	Aquino – 50g/L Fenpicoxamid (Inatreq) (FRAC Code 21) Turret 60 – 60g/L Metconazole (FRAC Code 3)		
Pack Size:	1 x 7.5L + 1 x 5L		
Formulation:	Aquino - Emulsifiable Concentrate Turret 60 – Emulsifiable Concentrate		
Maximum Single Dose Rate:	Aquino - 2L/ha Turret 60 – 1.5L/ha		
Maximum Total Dose per crop:	Aquino - 2L/ha Turret 60 – 3L/ha		
Maximum No. of Applications:	1 per crop per season		
Application Timing:	Beginning of stem elongation (GS 30)- end of flowering (GS 69)		
Water Volume:	100-300L/ha. But ideally use a minimum spray volume of 200l/ha in line with Corteva's 2023 Inatreq application advice. Spray pressure at 2-3 bar is recommended		
Aquatic buffer zone	Buffer zone is determined by the STRIPE Water Tool depending on rate and nozzle type		

Best Use Advice

- Aquino may be used on all commercial varieties of winter crops of wheat, triticale, durum wheat and spelt wheat.
- Ideally apply at T2 to see maximum yield benefits but Aquino could alternatively be applied at T1 if preferred.
- Apply at the recommended rate of 2L/ha.
- Aquino will give the best results before disease has become established in the crop and offers outstanding control of *Septoria* both with persistent protectant and curative activity.
- Aquino offers broad spectrum, superior disease control which leads to longer green leaf retention and improved yield.
- Rainfastness is 1 hour for Aquino.

Resistance Management

- Aquino MUST always be used in a mixture with a partner product which has a different mode of action. Available in an Ag pack with Turret 60, these products work together for complete control.
- Turret 60 adds to Aquino's control of yellow and brown rust.
- You must not apply more than one application of Aguino to any cereal crop in the same year.
- Aquino should be used in accordance with the instructions for use for the target disease at the specific growth rate indicated.



Tank Mixing and Application

- Aquino + Turret 60 has been tested in tank mixes with a variety of PGRs, herbicides, trace elements and other key fungicides.
- Products should only be tank mixed if each product can be applied with the label recommendations for its use.
- For a full tank mix list please visit the website- https://www.corteva.ie/tools-and-advice/tank-mixes.html
- Before applying Aquino + Turret 60, please see our best practice application advice and resources to help optimise field operations.
 Visit www.corteva.ie/inatregadvice or scan the QR code.

Disease Spectrum

Where applied with Turret 60, Aquino will give control to the following diseases in the following crops shown in the table below.

Disease	Winter Wheat	Winter Durum Wheat	Winter Triticale	Winter Spelt Wheat
Septoria Leaf Blotch	С	С	С	С
Yellow Rust	С	С	С	С
Brown Rust	С	С	С	С
				C=Control

STRIPE- Surface Water Tool for Reducing the Impact of Pesticides in the Environment

STRIPE is an initiative which allows farmers to reduce the size of mandatory untreated areas of land near water courses (buffer zones) while helping to protect aquatic life from pesticide contamination by reducing exposure.

Buffer zones are determined by the STRIPE water tool when ditches are present.

The buffer zone may be reduced by the use of low drift nozzles and application rate as per the following table.

	Full Rate	¾ Rate	½ Rate
Non- Drift reducing nozzles	30m	23m	15m
50% Drift reducing nozzles	15m	11m	7m
75% Drift reducing nozzles	7m	6m	4m
90% Drift reducing nozzles	3m	2m	1m

